arXiv:1404.3480v1 [math.NT] 14 Apr 2014

On a Lower Bound for $\|(4/3)^k\|$

Yury PUPYREV

April 15, 2014

Abstract

We prove, that

$$\left\|\left(\frac{4}{3}\right)^k\right\| > \left(\frac{4}{9}\right)^k \qquad \text{for} \quad k \geq 6,$$

where $\|\cdot\|$ is a distance to the nearest prime.

1 Introduction

In 1994 Bennett [1] considered a generalization of Waring's problem, namely, a problem on the order $g_N(k)$ of the additive basis

$$S_N^{(k)} = \{1^k, N^k, (N+1)^k, \dots\}, \qquad N \geq 2,$$

of the set of positive integers. He established the following estimates for $\|(1+1/N)^k\|$:

$$\left\|\left(1+\frac{1}{N}\right)^k\right\| > 3^{-k} \qquad \text{for} \quad 4 \leq N \leq k \cdot 3^k$$

and with their help obtained the representation

$$g_N(k) = N^k + \left[\left(1+\frac{1}{N}\right)^k\right] - 2$$

for $4 \leq N \leq (k+1)^{(k-1)/k} - 1$.

He concluded that he needed the inequality

$$\left\|\left(\frac{4}{3}\right)^k\right\| > \left(\frac{4}{9}\right)^k \qquad \text{for} \quad k \geq 6, \tag{1}$$

for the representation

$$g_3(k) = 3^k + \left[\left(\frac{4}{3}\right)^k\right] - 2.$$

In 2007 Zudilin [6], by modifying Baker's construction, namely, by considering Padé approximations to the remainder of the series

$$\frac{1}{(1-z)^{m+1}} = \sum_{n=0}^{\infty} \binom{m+n}{m} z^n$$

and by receiving sharp estimates for the p-adic orders of the arising binomial coefficients, arrived at the bound

$$\left\| \left(\frac{4}{3}\right)^k \right\| > 0.4914^k \qquad \text{for} \quad k \ge K,$$

where K is an effective constant.

In 2009 this author [4] received an exact value of K, but it was too big for checking (1) for $6 \le k < K$.

In this paper using the same method as Zudilin, but with another set of parameters, we receive the bound (1) for $k \ge 17\,545\,718$, and check it using a lemma similar to [2, Proposition 1], with software for remaining k.

Thus, we prove the following result.

Theorem 1. *We have*

$$\left\| \left(\frac{4}{3}\right)^k \right\| > \left(\frac{4}{9}\right)^k \qquad \textit{holds for} \quad k \ge 6.$$

2 Padé Approximations

Following [6], we fix two positive integers a and b, with $3a \le b$, and write

$$\begin{aligned}
\left(\frac{4}{3}\right)^{2(b+1)} &= \left(\frac{16}{9}\right)^{(b+1)} = 2^{b+1}\left(1+\frac{1}{8}\right)^{-(b+1)} \\
&= 2^{b+1} \sum_{l=0}^{\infty} \binom{b+l}{b} \left(\frac{1}{8}\right)^l (-1)^l \\
&= 2^{b+1} 2^{-3a} \sum_{l=0}^{\infty} \binom{b+l}{b} 2^{3(a-l)} (-1)^{-l} \\
&= 2^{b-3a+1} (-1)^a \sum_{l=0}^{\infty} \binom{b+l}{b} 2^{3(a-l)} (-1)^{(a-l)} \\
&\equiv 2^{b-3a+1} (-1)^a \sum_{l=a}^{\infty} \binom{b+l}{b} 2^{3(a-l)} (-1)^{(a-l)} \pmod{\mathbb{Z}} \\
&\equiv 2^{b-3a+1} (-1)^a \sum_{\nu=0}^{\infty} \binom{a+b+\nu}{b} \left(-\frac{1}{8}\right)^{\nu} \pmod{\mathbb{Z}}. \qquad (2)
\end{aligned}$$

So, we are going to consider Padé approximations to the function

$$F(z) = F(a,b;z) = \sum_{\nu=0}^{\infty} \binom{a+b+\nu}{b} z^{\nu}. \tag{3}$$

For any positive integer $n \le b$ we find [6]

$$Q_n(z^{-1}) = \frac{(a+b+n)!}{(a+n-1)!\,n!\,(b-n)!} \int_0^1 t^{a+n-1}(1-t)^{b-n}(1-z^{-1}t)^n\,dt \tag{4}$$

and

$$\begin{aligned} R_n(z) = \frac{(a+b+n)!}{(a+n-1)!\,n!\,(b-n)!} \cdot z^n \\ \times \int_0^1 t^n(1-t)^{a+n-1}(1-zt)^{-(a+b+n)-1}\,dt \end{aligned} \tag{5}$$

such that

$$Q_n(z^{-1})F(z) = P_n(z^{-1}) + R_n(z), \tag{6}$$

is performed with polynomial $P_n(x) \in \mathbb{Z}[x]$, $\deg P_n \le n-1$.

3 Arithmetic argument

For every prime $p > \sqrt{a+b+n}$ we set

$$\begin{aligned} e_{p,n} = \min_{\mu\in\mathbb{Z}}\Bigg(-\left\{-\frac{a+n}{p}\right\} + \left\{-\frac{a+n+\mu}{p}\right\} + \left\{\frac{\mu}{p}\right\} \\ - \left\{\frac{a+b+n}{p}\right\} + \left\{\frac{a+b+\mu}{p}\right\} + \left\{\frac{n-\mu}{p}\right\}\Bigg), \\ e'_{p,n} = \min_{\mu\in\mathbb{Z}}\Bigg(-\left\{\frac{a+n+\mu}{p}\right\} + \left\{\frac{a+n}{p}\right\} + \left\{\frac{\mu}{p}\right\} \\ - \left\{\frac{a+b+n}{p}\right\} + \left\{\frac{a+b+\mu}{p}\right\} + \left\{\frac{n-\mu}{p}\right\}\Bigg), \end{aligned}$$

and for

$$\Phi = \Phi(a,b,n) = \prod_{p>\sqrt{a+b+n}} p^{e_{p,n}}, \qquad \Phi' = \Phi'(a,b,n) = \prod_{p>\sqrt{a+b+n}} p^{e'_{p,n}},$$

by lemmas 3 and 4 in [6], we have

$$\begin{aligned} \Phi^{-1}Q_n(x),\ \Phi^{-1}P_n(x) \in \mathbb{Z}[x], \\ \Phi'^{-1}(n+1)Q_{n+1}(x),\ \Phi'^{-1}(n+1)P_{n+1}(x) \in \mathbb{Z}[x]. \end{aligned}$$

4 A Bound for $\|(4/3)^k\|$

For a, b, and n we write

$$a = \alpha m, \quad b = \beta m, \quad n = \gamma m, \qquad m \in \mathbb{N}.$$

Our aim is to find a lower bound for the absolute value of ε_k, where

$$\left(\frac{4}{3}\right)^k = M_k + \varepsilon_k, \qquad M_k \in \mathbb{Z}, \quad 0 < |\varepsilon_k| < \frac{1}{2}.$$

For $k \geq 3$ we write $k = 2(\beta m + 1) + j$ with positive integers m and $j < 2\beta$. We multiply both sides of (6) by $\widetilde{\Phi}^{-1} 2^{b-3a+1+2j}(-1)^a$ (where $\widetilde{\Phi}$ is equal to Φ or to $\Phi'/(n+1)$; we discuss this choice in what follows) and put $z = -1/8$:

$$\begin{aligned} &Q_n(-8)\widetilde{\Phi}^{-1}3^j \cdot \left(\frac{4}{3}\right)^j 2^{b-3a+1}(-1)^a F\left(a, b, -\frac{1}{8}\right) \\ &\quad = P_n(-8)\widetilde{\Phi}^{-1}2^{b-3a+1+2j}(-1)^a + R_n\left(-\frac{1}{8}\right)\widetilde{\Phi}^{-1}2^{b-3a+1+2j}(-1)^a. \end{aligned} \tag{7}$$

From (2) and (3) one can find that

$$\left(\frac{4}{3}\right)^j 2^{b-3a+1}(-1)^a F\left(a, b, -\frac{1}{8}\right) \equiv \left(\frac{4}{3}\right)^{2(b+1)+j} \pmod{\mathbb{Z}} = \left(\frac{4}{3}\right)^k,$$

so the left-hand side can be written as $M'_k + \varepsilon_k$ and one can rewrite (7) as

$$Q_n(-8)\widetilde{\Phi}^{-1}3^j \cdot \varepsilon_k = M''_k + R_n\left(-\frac{1}{8}\right)\widetilde{\Phi}^{-1}2^{b-3a+1+2j}(-1)^a. \tag{8}$$

At this point we should check if the number M''_k is distinct from zero. Lemma 2 in [6] guarantees that for n or for $n+1$ we have $M''_k \neq 0$. So $\widetilde{\Phi} = \Phi$ if (8) holds for n with $M''_k \neq 0$, and $\widetilde{\Phi} = \Phi'/(n+1)$ if (8) holds for $n+1$ with $M''_k \neq 0$. (This odd way of working with $\widetilde{\Phi}$ becomes more understandable once we have determined a, b, and n.)

So, assuming that

$$\left| R_n\left(-\frac{1}{8}\right)\widetilde{\Phi}^{-1}2^{b-3a+1+2j}(-1)^a \right| < \frac{2}{3}, \tag{9}$$

from (8) we have

$$|Q_n(-8)\widetilde{\Phi}^{-1}3^j| \cdot |\varepsilon_k| \geq |M''_k| - \left| R_n\left(-\frac{1}{8}\right)\widetilde{\Phi}^{-1}2^{b-3a+1+2j}(-1)^a \right| > \frac{1}{3},$$

and so

$$|\varepsilon_k| > \frac{\widetilde{\Phi}}{3^{j+1}|Q_n(-8)|} \geq \frac{\widetilde{\Phi}}{3^{2\beta}|Q_n(-8)|},$$

which means that

$$|\varepsilon_k| > \frac{\Phi}{3^{2\beta}|Q_n(-8)|}\,. \tag{10}$$

or

$$|\varepsilon_k| > \frac{\Phi'}{(n+1)3^{2\beta}|Q_{n+1}(-8)|}\,, \tag{11}$$

depending on the choice made in (8).

5 A Bound for Φ

For evaluating Φ and Φ' we consider the functions

$$\begin{aligned}\varphi(x) = \min_{0\le y<1}(&-\{-(\alpha+\gamma)x\} + \{-(\alpha+\gamma)x - y\} + \{y\}\\ &- \{(\alpha+\beta+\gamma)x\} + \{(\alpha+\beta)x + y\} + \{\gamma x - y\}),\\ \varphi'(x) = \min_{0\le y<1}(&-\{(\alpha+\gamma)x + y\} + \{(\alpha+\gamma)x\} + \{y\}\\ &- \{(\alpha+\beta+\gamma)x\} + \{(\alpha+\beta)x + y\} + \{\gamma x - y\}),\end{aligned}$$

which take the values $e_{n,p}$ and $e'_{n,p}$, respectively at the point m/p.

All the solutions x of the equation $\varphi(x) = 1$ form the set of intervals in $[0, 1)$, which should contain $\{x\}$. If we denote A_i and B_i the left and right points of this intervals, respectively, then the condition $A_i \le \{m/p\} < B_i$ (i.e. $e_{n,p} = \varphi(m/p) = 1$) is equivalent to

$$A_i + N \le \frac{m}{p} < B_i + N, \qquad N \in \mathbb{N},$$

($\mathbb{N}$ is the set of non-negative integers), or the same

$$p \in \left(\frac{m}{B_i + N}\,, \frac{m}{A_i + N}\right], \qquad N \in \mathbb{N}.$$

This means that all the prime numbers p such that

$$p \in \bigcup_{N=0}^{\left[\frac{m}{\sqrt{a+b+n}}\right]-1} \bigcup_i \left(\frac{m}{B_i + N}\,, \frac{m}{A_i + N}\right]$$

(the inequality $p > \sqrt{a+b+n}$ entails $m/(B_i + N) \ge \sqrt{a+b+n}$, and one can find the bound for N) go to Φ. So we have

$$\log\Phi \ge \sum_{N=0}^{\left[\frac{m}{\sqrt{a+b+n}}\right]-1} \sum_i \left(\theta\left(\frac{m}{A_i + N}\right) - \theta\left(\frac{m}{B_i + N}\right)\right), \tag{12}$$

where $\theta(x) = \sum_{p\le x,\, p \text{ is prime}} \log p$.

The same works for $\varphi'(x)$. And it is proved in [6], that the sets for $\varphi(x)$ and $\varphi'(x)$ differs only on a set of zero measure.

6 Analytic and Arithmetical Bounds

Let us take

$$\alpha = 3, \qquad \beta = 9, \qquad \gamma = 4.$$

For Φ we have the set of intervals

$$\left[\frac{1}{8}, \frac{1}{7}\right] \cup \left[\frac{3}{16}, \frac{1}{5}\right) \cup \left[\frac{3}{8}, \frac{2}{5}\right) \cup \left[\frac{9}{16}, \frac{4}{7}\right] \cup \left[\frac{11}{16}, \frac{5}{7}\right] \cup \left[\frac{15}{16}, 1\right)$$

For Φ' the difference will only be in the right-end points of the intervals.

We will use the following bounds for $\theta(x)$ [5]: the upper bound

$$\theta(x) < 1.001102 \cdot x \qquad \text{if} \quad x > 0,$$

and the lower bound

$$\theta(x) > 0.998 \cdot x \qquad \text{if} \quad x > 487\,381.$$

Substituting them in (12) and taking the sum for $N = 0, 1$ we obtain

$$\log \Phi > 1.639533 \cdot m \qquad \text{for} \quad m \geq 974\,762. \tag{13}$$

The same bound holds for Φ'.

We need to estimate the values of $|Q_n(z^{-1})|$, $|R_n(z)|$, and $|Q_{n+1}(z^{-1})|$, $|R_{n+1}(z)|$ at the point $z = -1/8$, and to estimate Φ'. We begin with

$$\log\left(\frac{(16m)!}{(7m-1)!\,(4m)!\,(5m)!}\right) \qquad \text{and} \qquad \log\left(\frac{(16m+1)!}{(7m)!\,(4m+1)!\,(5m-1)!}\right).$$

We use Stirling's formulae [3]

$$\sqrt{2\pi n}\left(\frac{n}{e}\right)^n < n! < \sqrt{2\pi n}\left(\frac{n}{e}\right)^n e^{1/(12n)},$$

and we find

$$\begin{aligned}
\log\left(\frac{(16m)!}{(7m-1)!\,(4m)!\,(5m)!}\right) &= \log\left(\frac{(16m)!}{(7m)!\,(4m)!\,(5m)!}\right) + \log(7m) \\
&< \frac{1}{2}\log(2\pi) + \frac{1}{2}\log(16) + \frac{1}{2}\log(m) + 16m\log(16) + 16m\log(m) - 16m \\
&\quad + \frac{1}{12 \cdot 16m} \\
&\quad - \left(\frac{1}{2}\log(2\pi) + \frac{1}{2}\log(7) + \frac{1}{2}\log(m) + 7m\log(7) + 7m\log(m) - 7m\right) \\
&\quad - \left(\frac{1}{2}\log(2\pi) + \frac{1}{2}\log(4) + \frac{1}{2}\log(m) + 4m\log(4) + 4m\log(m) - 4m\right) \\
&\quad - \left(\frac{1}{2}\log(2\pi) + \frac{1}{2}\log(5) + \frac{1}{2}\log(m) + 5m\log(5) + 5m\log(m) - 5m\right) \\
&\quad + \log(7) + \log(m).
\end{aligned}$$

Since

$$\begin{aligned}
&-\log(2\pi)+\frac{1}{2}(\log(16)-\log(7)-\log(4)-\log(5))-\log(m)\\
&\qquad+\frac{1}{12\cdot 16m}+\log(7)+\log(m)\\
&\quad<0 \qquad \text{for} \quad m\geq 974\,762.
\end{aligned}$$

one can have

$$\begin{aligned}
&\log\left(\frac{(16m)!}{(7m-1)!\,(4m)!\,(5m)!}\right)\\
&\quad<(16\log(16)-7\log(7)-4\log(4)-5\log(5))\cdot m \qquad (14)\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{for} \quad m\geq 974\,762.
\end{aligned}$$

Now,

$$\begin{aligned}
&\log\left(\frac{(16m+1)!}{(7m)!\,(4m+1)!\,(5m-1)!}\right)\\
&\quad=\log\left(\frac{(16m)!}{(7m)!\,(4m)!\,(5m)!}\right)+\log\left(\frac{(16m+1)(5m)}{(4m+1)}\right)\\
&\quad<\log\left(\frac{(16m)!}{(7m)!\,(4m)!\,(5m)!}\right)+\log\left(\frac{(16m+4)(5m)}{(4m+1)}\right)\\
&\quad=\log\left(\frac{(16m)!}{(7m)!\,(4m)!\,(5m)!}\right)+\log(20m),
\end{aligned}$$

and in a similar way we conclude that

$$\begin{aligned}
&\log\left(\frac{(16m+1)!}{(7m)!\,(4m+1)!\,(5m-1)!}\right)\\
&\quad<(16\log(16)-7\log(7)-4\log(4)-5\log(5))\cdot m+1 \qquad (15)\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{for} \quad m\geq 974\,762.
\end{aligned}$$

For the integral in (5) we write the estimates

$$\begin{aligned}
&\int_0^1 t^{4m}(1-t)^{7m-1}(1-zt)^{-16m-1}\,dt\\
&\qquad<\left(\max_{t\in[0,1]} t^4(1-t)^7\left(1+\frac{t}{8}\right)^{-16}\right)^{m-1}\int_0^1 t^4(1-t)^6\left(1+\frac{t}{8}\right)^{-17}dt,\\
&\int_0^1 t^{4m+1}(1-t)^{7m}(1-zt)^{-16m-2}\,dt\\
&\qquad<\left(\max_{t\in[0,1]} t^4(1-t)^7\left(1+\frac{t}{8}\right)^{-16}\right)^{m-1}\int_0^1 t^5(1-t)^7\left(1+\frac{t}{8}\right)^{-18}dt,
\end{aligned}$$

and so

$$\log\left(\int_0^1 t^{4m}(1-t)^{7m-1}(1-zt)^{-16m-1}\,dt\right) < -7.884160\cdot(m-1)-8.568400,$$

$$\log\left(\int_0^1 t^{4m+1}(1-t)^{7m}(1-zt)^{-16m-2}\,dt\right) < -7.884160\cdot(m-1)-10.140038.$$

Let us check inequality (9) for n and $n+1$:

$$\begin{aligned}
&\log\left|R_n\left(-\frac{1}{8}\right)\widetilde{\Phi}^{-1}2^{b-3a+1+2j}(-1)^a\right| \\
&\quad< 17.147682\cdot m+\log\left(\frac{1}{8}\right)^4\cdot m-7.884160\cdot(m-1)-8.568400 \\
&\qquad-1.639533\cdot m+35\log(2) \\
&\quad< -0.693777\cdot m+23.575912<\log\left(\frac{2}{3}\right)\qquad\text{for}\quad m\ge 974\,762,\\
&\log\left|R_{n+1}\left(-\frac{1}{8}\right)\widetilde{\Phi}^{-1}2^{b-3a+1+2j}(-1)^a\right| \\
&\quad< 17.147682\cdot m+1+\log\left(\frac{1}{8}\right)^4\cdot m+\log\left(\frac{1}{8}\right)-7.884160\cdot(m-1) \\
&\qquad-10.140038-1.639533\cdot m+\log(4)\log(m)+1+35\log(2) \\
&\quad< -0.693777\cdot m+\log(4)\log(m)+21.924832<\log\left(\frac{2}{3}\right) \\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\text{for}\quad m\ge 974\,762.
\end{aligned}$$

So, inequality (9) holds, and we can move on.

For integral in (4), in the same way as for the one in (5), one has

$$\log\left(\int_0^1 t^{7m-1}(1-t)^{5m}(1+8t)^{4m}\,dt\right) < -0.945755\cdot(m-1)-1.725707,$$

$$\log\left(\int_0^1 t^{7m}(1-t)^{5m-1}(1+8t)^{4m+1}\,dt\right) < -0.945755\cdot(m-1)+0.878883.$$

Now we can calculate the bounds in (10) and (11). We begin with (10):

$$\begin{aligned}\log|\varepsilon_k| &> 1.639533\cdot m - 18\log(3) - 17.147682\cdot m \\ &\quad + 0.945755\cdot(m-1) + 1.725707 \\ &> -14.562394\cdot m - 18.995070 \\ &> -0.81\cdot k > \log\left(\frac{4}{9}\right)\cdot k \qquad \text{for} \quad k \geq 17\,545\,718.\end{aligned}$$

For (11) we have

$$\begin{aligned}\log|\varepsilon_k| &> 1.639533\cdot m - \log(4m+1) - 18\log(3) - 17.147682\cdot m \\ &\quad + 0.945755\cdot(m-1) - 0.878883 \\ &> -14.562394\cdot m - \log(4)\log(m) - 22.599660 \\ &> -14.562414\cdot m - 22.599660 \\ &> -0.81\cdot k > \log\left(\frac{4}{9}\right)\cdot k \qquad \text{for} \quad k \geq 17\,545\,718.\end{aligned}$$

So, we have

$$\left\|\left(\frac{4}{3}\right)\right\| > \left(\frac{4}{9}\right)^k \qquad \text{for} \quad k \geq 17\,545\,718.$$

7 The Final check

We need to check inequality (1) for $6 \leq k \leq 17\,545\,717$. Following [2], we prove next lemma.

Lemma 1. *Let m be a positive integer, and assume that the number 4^m contain no block of h consecutive 0, or 2, in its ternary expansion. Then the inequality*

$$\left\|\left(\frac{4}{3}\right)^k\right\| \geq \left(\frac{4}{9}\right)^k \tag{16}$$

holds for all

$$m\left(\frac{\log 4}{\log 9}\right) + \frac{h}{2} \leq k \leq m. \tag{17}$$

Proof. We give a proof by contradiction. Assume that k is in the specified interval, but (16) is not true. Then for some integer M_1 we have one of the next two equalities:

$$M_1 = \left(\frac{4}{3}\right)^k \pm \epsilon_1, \qquad \text{where} \quad 0 < \epsilon_1 < \left(\frac{4}{9}\right)^k,$$

so, with some integer M_2

$$4^m = 3^k M_2 \mp \epsilon_2, \qquad \text{where} \quad 0 < \epsilon_2 < 4^m 3^{-k}.$$

Since $m(\log 4/\log 9) \le k - h/2$, we have

$$0 < \epsilon_2 < 9^{k-h/2} \cdot 3^{-k} = 3^{k-h},$$

but this means, that a block of h digits of the number 4^m, which are responsible for powers $3^{k-h}, 3^{k-h+1}, \ldots, 3^{k-1}$, consists of 2, or 0. □

For specified m the software calculates $h(m)$ defined in lemma 1, descends to the new value of m prescribed by (17), and so on. We started calculations with $m = 17\,545\,718$ and stopped at $m = 5$. Results of all the steps are given in Table 1.

Table 1:

	m	h
1	17545718	18
2	11229269	16
3	7186741	16
4	4599523	14
5	2943702	14
6	1883977	14
7	1205753	15
8	771689	11
9	493886	13
10	316094	12
11	202307	11
12	129482	10
13	82874	12

	m	h
14	53046	11
15	33955	10
16	21737	11
17	13917	8
18	8911	9
19	5708	8
20	3658	10
21	2347	8
22	1507	7
23	968	5
24	622	7
25	402	10
26	263	5

	m	h
27	171	4
28	112	4
29	74	3
30	49	3
31	33	3
32	23	3
33	16	3
34	12	3
35	9	2
36	7	1
37	5	2

So Theorem 1 is proved.

The author expresses his gratitude to Igor P. Rochev for his help in preparing the paper.

References

[1] M. A. Bennett, *An ideal Waring problem with restricted summands*. Acta Arith. **66**:2, (1994), 125–132.

[2] F. Delmer, J.-M. Deshouillers, *The computation of $g(k)$ in Waring's problem*. Math. Comp. **54** (1990), 885–893.

[3] G. M. Fikhtengolts, *The Course of Differential and Integral Calculus*, Vol. 2, Nauka, Moscow, 1966 (in Russian).

[4] Yu. A. Pupyrev, *Effectivization of a lower bound for $\|(4/3)^k\|$*. Math. Notes **85** (2009), 877–885.

[5] J. B. Rosser, L. Schoenfeld, *Sharper Bounds for the Chebyshev Functions $\theta(x)$ and $\psi(x)$*. Math. Comp. **29** (1975), 243–269.

[6] W. Zudilin, *A new lower bound for $\|(3/2)^k\|$*. J. Théor. Nombres Bordeaux **19** (2007), 311–323